

# MEDICAL SCIENCE

## To Cite:

Agarwal A, Hamayun R, Fahmy EK, Alanazi KS, Alanazi AMM, Makhdoom AK, Alanazi AMM. Prevalence and awareness of hypertension among the shopkeepers working in the market in Northern Border Province of Saudi Arabia. *Medical Science* 2023; 27: e376ms3237 doi: <https://doi.org/10.54905/disssi.v27i141.e376ms3237>

## Authors' Affiliation:

<sup>1</sup>Department of Pathology, Faculty of Medicine, Northern Border University, Arar, KSA

<sup>2</sup>Department of Pharmacology, Faculty of Medicine, Northern Border University, Arar, KSA

<sup>3</sup>Department of Physiology, Faculty of Medicine, Northern Border University, Arar, KSA

<sup>4</sup>MBBS Students, Faculty of Medicine, Northern Border University, Arar, KSA

## \*Corresponding author

MBBS Students, Faculty of Medicine, Northern Border University, Arar, KSA

Email: [akm.nbu@gmail.com](mailto:akm.nbu@gmail.com)

## ORCID List

Anshoo Agarwal	0000-0002-7435-0404
Rahma Hamayun	0000-0002-7731-5065
Eslam Kamal Fahmy	0000-0002-1994-2570
Khalid Saud Alanazi	0000-0002-5169-2524
Abdulaziz Mohammed M Alanazi	0000-0002-0255-5945
Amin Khalid Makhdoom	0000-0001-5434-6988
Abdulaziz Mohammed Mukhleef Alanazi	0000-0001-6409-7153

## Peer-Review History

Received: 21 August 2023

Reviewed & Revised: 25/August/2023 to 03/November/2023

Accepted: 07 November 2023

Published: 11 November 2023

## Peer-review Method

External peer-review was done through double-blind method.

Medical Science

pISSN 2321-7359; eISSN 2321-7367



© The Author(s) 2023. Open Access. This article is licensed under a [Creative Commons Attribution License 4.0 \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



# Prevalence and awareness of hypertension among the shopkeepers working in the market in Northern Border Province of Saudi Arabia

Anshoo Agarwal<sup>1</sup>, Rahma Hamayun<sup>2</sup>, Eslam Kamal Fahmy<sup>3</sup>, Khalid Saud Alanazi<sup>4</sup>, Abdulaziz Mohammed M Alanazi<sup>4</sup>, Amin Khalid Makhdoom<sup>4\*</sup>, Abdulaziz Mohammed Mukhleef Alanazi<sup>4</sup>

## ABSTRACT

**Background:** The public health systems of developing and developed countries are seriously threatened by hypertension because it is a widespread non-communicable disease and a global pandemic. We anticipate a high incidence of cardiovascular disease risk factors in our study because we assume that shopkeepers within the market population lead sedentary lifestyles as most of their jobs involve not much physical activity. **Aim and objectives:** To know the prevalence and awareness of risk factors for cardiovascular disease (CVD) and the relation with the dietary patterns in the market population in Arar, Kingdom of Saudi Arabia (KSA). **Material & methods:** In a randomized cross-sectional study, a population of market workers was surveyed for cardiovascular diseases and their risk factors by giving a pre-validated structured questionnaire. **Results:** The prevalence of hypertension and obesity was found to be high in the market population of Arar. Hypertension was more prevalent in overweight and obese individuals. A sedentary lifestyle among shopkeepers who continue to sit for long hours was more prevalent in obese and in the pre-hypertensive or pre-diabetic category. **Conclusions:** A wholesome diet awareness corresponds with knowledge of high-risk factors for cardiovascular diseases (CVD), corresponding to the hypothesized link between nutritional knowledge and diet quality. It may influence poor dietary behavior found among the market population, and it may help reduce the incidence of hypertension, heart attack, stroke, and other associated risk factors for CVD.

**Keywords:** Prevalence, CVD, Hypertension, Market Population, Saudi Arabia

## 1. INTRODUCTION

Since hypertension is a common non-communicable disease and a global pandemic, it poses a severe threat to both emerging and industrialized public health infrastructures (Roger et al., 2012). The prevalence of cardiovascular diseases has been rising on a worldwide basis. Age, gender, ethnicity, and physical attributes of an individual significantly impact the prevalence rates of cardiovascular diseases. Cardiovascular diseases are among the leading causes of mortality in Saudi Arabia. Cardiovascular diseases, cerebrovascular disorders, blindness, and kidney diseases are associated with uncontrolled blood pressure (Ng et al., 2018). Proper blood pressure control can help to avoid severe complications of hypertension.

We expect a high incidence of cardiovascular diseases in our research group of the market population since we assume they live sedentary lifestyles because most of their work requires them to sit in a single place (Padmanabhan et al., 2018). Besides living a sedentary lifestyle, most of them don't regularly exercise. The workers would spend most of their day at the market, relying on food vendors and eating fast food for most of their meals (Roy et al., 2017). Foods rich in oil and salt make them a risk factor for cardiovascular diseases. For awareness among the market population, extensive health education addressing the risk factors for cardiovascular diseases was needed (Forouzanfar et al., 2017).

Appropriate hypertension control is necessary to lower the mortality rate linked to hypertension, the most prevalent CVD risk factor (Al-Nozha et al., 1997). Hypertension is linked to 23% of all deaths and 32% of adult deaths (Al-Hamdan et al., 2011). Numerous studies have found a connection between high blood pressure and millions of disability-adjusted life years (DALYs) (Saeed et al., 2011). The load of hypertension expected to increase significantly in the forthcoming years due to stressful environmental and lifestyle changes that stem from dangerous working conditions and growing societal pressures (Memish et al., 2014).

### Aims and objectives

To know the prevalence and awareness of risk factors for cardiovascular disease (CVD) in the market population in Arar, Kingdom of Saudi Arabia (KSA)

To study the dietary pattern of the market population in Arar, Kingdom of Saudi Arabia.

To analyze the relationship of diet patterns with prevalence risk factors for cardiovascular disease in the market population in Arar, Kingdom of Saudi Arabia (KSA)

## 2. MATERIALS AND METHODS

### Study Design and Participants

In a randomized cross-sectional study, cardiovascular diseases and their related risk factors were surveyed in the market population. To check the demographic details and awareness regarding cardiovascular diseases, a structured questionnaire was distributed to the market population in Arar, Saudi Arabia. The sample included 333 shopkeepers and workers in the market in Arar, Saudi Arabia.

### Research tool

A pre-structured questionnaire designed after a literature review. The dietary pattern of the market population in Arar, Saudi Arabia, was studied using a food frequency questionnaire (Geaney et al., 2015). All participants were given a self-administered questionnaire (in both English and Arabic). A survey conducted for the enrolment of the participants. The consent obtained from the willing participants. Participants filled a pre-validated survey questionnaire.

### Study tool

The survey questionnaire contained two sections. The first section included items related to the demographic information of the participants. The second section had information to assess the Food frequency analysis of participants.

### Study setting

Markets of Arar City, Saudi Arabia, from 1/1/2023 to 1/2/2023.

### Study population

Market worker population in Arar City.

**Study Design**

Cross-sectional study.

**Sampling tool**

Structured, well-designed questionnaire (both English and Arabic versions).

**Sampling Method**

Simple random sample

**Sampling technique**

Simple random sample

**Sample size**

333 Market Population workers

The sample size was calculated using the formula  $n = \frac{z^2 p (1-p)}{d^2}$  where  $p$ =prevalence found in the previous study is 0.22,  $d$ =tolerated error is 0.05, Confidence interval=95%, and  $Z$  score =1.96.

**Inclusion and Exclusion criteria*****Inclusion criteria***

Males and females working in the market in Arar.

***Exclusion criteria***

Pregnant females and children and the elderly population.

Males and females who had been undergoing treatment for any chronic illnesses

**Data Analysis*****Statistics***

The data obtained from the questionnaire analysis was entered into Microsoft Excel 2013. Microsoft Excel and Statistical Package for Social Science (SPSS) software program (IBM SPSS v.20 Inc., Chicago IL, USA) were used for descriptive analysis of the data. The results were presented as frequency distribution.  $P$ -value< 0.05 was considered significant.

### 3. RESULTS

The socio-demographic details of the participants ( $n = 333$ ) given in (Table 1). The average height was 158 cm, and the average weight was 82 kg. The majority of the people had poor dietary habits, and they were unaware that nutritional habits and physical fitness are related to cardiovascular diseases and their risk factors. The food habits of participants were assessed using the food frequency questionnaire listed in (Tables 2-6) and (Figure 1). The prevalence of hypertension and obesity, which are major risk factors for cardiovascular diseases, were found to be high in the market population of Arar. A sedentary lifestyle (39%) among shopkeepers who continue to sit for long hours were found to be more obese and in pre-hypertensive or pre-diabetic people.

**Daily Physical Activity Information**

52% of participants stated they had been active in general, and 45% had no weekly physical activity.

**Blood pressure treatment information**

27% of participants had taken medicine to lower blood pressure.

**History of Diabetes-mellitus-related information**

23% of participants had been taking diabetes-mellitus treatment, 18% had been doing Self-Monitoring of Blood Glucose, 27% had a family history of diabetes-mellitus, 39% had a family history of heart disease and 59% stated that they are under stress all the time.

Results of Food Frequency Questionnaire-FFQ (Total number of participants=333)

The FFQ approved for usage in a variety of populations. Participants tracked how frequently they typically ate each meal item throughout one month. The FFQ includes 150 food items divided into the major food groups and evaluated the entire diet. The frequency of consumption of each food item was recorded and converted into amounts using typical serving sizes. Information collected was translated to food quantities and nutrient values using NetWISP4 (Weighed Intake Software application; Tinuviel Software, Warrington, UK), a specially developed nutrition software application. Some 333 participants were involved in this study. 52.9% were single, and 40.2% were married. The majority have a university educational background, 72.4%. Smokers in the study were 57.4%. Having a first-degree family member with hypertension was 43.2%. And more than half of the participants, 52.9%, had standard body mass index (Table 1).

Table 1 Socio-demographic characteristics

Characteristics		N (%)	Total
Social status	Single	52.9%	333
	Married	40.2%	
	Divorced	6.9%	
Educational background	High school	27.6%	
	University	72.4%	
Do you smoke	Yes	42.6%	
	No	57.4%	
Family history (first-degree) of hypertension	Yes	43.2%	
	No	36.6%	
	I do not	20.1%	
Body mass index categories	Underweight	11.1%	
	Normal body mass index	52.9%	
	Overweight	25.8%	
	Obese	10.2%	




Figure 1 Meat consumption per month

40.4% and 47.6% include burgers and fish 1-3 per month in their diet, respectively. Including Beef: Roast, steak, mince, and stew for Never or less than once/month, 1-3 per month, and 2-4 per week 29.2%, 29.2%, and 29.8%, respectively (Figure 1). 44.4% had low-fat yogurt never or less than once per month. 1-3 per month had Full-fat or Greek yogurt, Cottage cheese, low-fat soft cheese, Eggs as boiled, fried, and scrambled, and Salad cream, mayonnaise 31.8%, 28.2%, 23.4% and 25.2%, respectively (Table 2).

**Table 2** Dairy consumption per month

	Low-fat yogurt	Full-fat or Greek yogurt	Cottage cheese, low-fat soft cheese	Eggs as boiled, fried, scrambled	Salad cream, mayonnaise
Never or less than once/per month	44.4%	29.7%	40.8%	15.9%	52.9%
1-3 per month	27.6%	31.8%	28.2%	23.4%	25.2%
2-4 per week	20.7%	25.5%	23.7%	35.1%	15.9%
5-6 per week	3.9%	8.4%	3.9%	15.3%	3.0%
once per day	1.8%	2.7%	1.8%	6.0%	0.0%
2-3 per day	0.9%	0.9%	0.6%	1.5%	2.1%
4-5 per day	0.6%	0.6%	0.3%	1.2%	0.3%
More than 6 per day	0.0%	0.3%	0.6%	1.5%	0.6%

30.9% had sweet biscuits and chocolate in their diet 2-4 per week 35.1%, 29.7%, and 21.6% had cake, Buns, or pastries. Croissants, doughnuts, and Sugar were added to tea and coffee 1-3 per month, respectively. 41.1% had Ice cream and chocolate ices in their diet never or less than once per month (Table 3).

**Table 3** Snacks consumption per month

	Sweet biscuits, chocolate	Cake	Buns, pastries, e.g., croissants, doughnuts	Ice cream, chocolate ices	Sweets	Sugar added to tea, coffee	Peanuts or other nuts	Tomato ketchup	Pickles	Jam, honey
Never or less than once/per month	16.2%	22.5%	18.3%	41.1%	20.7%	19.8%	35.7%	34.8%	43.2%	41.7%
1-3 per month	20.1%	35.1%	29.7%	34.5%	26.7%	21.6%	30.3%	31.2%	28.2%	27.0%
2-4 per week	30.9%	30.9%	30.3%	18.9%	27.9%	27.6%	18.0%	19.5%	17.1%	18.6%
5-6 per week	18.6%	5.1%	11.7%	3.9%	11.1%	10.2%	10.5%	8.4%	5.1%	5.1%
once per day	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%
2-3 per day	3.0%	0.6%	0.6%	0.3%	1.5%	5.1%	1.8%	1.2%	0.3%	0.9%
4-5 per day	1.2%	0.3%	1.5%	0.6%	1.8%	3.0%	0.6%	0.9%	1.5%	0.9%
More than 6 per day	9.9%	5.4%	7.8%	0.6%	10.2%	12.6%	3.0%	3.9%	4.5%	5.7%

Participants consume tea, coffee, soft drinks, and pure fruit juice 2-4 per week 28.8%, 24.0%, 25.2%, and 27.3%. Apples, pears, bananas, grapes, and melon are the least consumed by the participants, never or less than one/month (Table 4).

**Table 4** Drinks and fruit consumption per month

	Tea	Coffee	Hot chocolate	Soft drinks	Pure fruit juice	Apple	Pears	Orange	Banana	Grapes	Melon	Peaches, plums, apricots	Strawberries, raspberries, kiwi fruit
Never or less than once/per month	13.8%	11.7%	40.8%	22.5%	25.8%	33.0%	42.6%	29.4%	30.3%	38.7%	45.9%	45.9%	42.6%
1-3 per month	13.2%	12.0%	28.8%	23.7%	34.2%	31.2%	30.3%	31.5%	30.9%	34.2%	29.7%	30.3%	31.2%
2-4 per week	28.8%	24.0%	15.3%	25.2%	27.3%	20.1%	16.5%	24.3%	22.8%	17.7%	15.3%	16.2%	17.1%
5-6 per week	12.9%	12.9%	6.6%	12.9%	5.4%	7.2%	3.9%	6.6%	6.9%	3.0%	4.5%	3.0%	3.0%
once per	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%

day													
2-3 per day	12.0%	15.3%	1.2%	4.2%	0.9%	1.8%	2.1%	2.1%	1.8%	2.4%	1.5%	1.2%	2.1%
4-5 per day	1.8%	3.3%	0.6%	1.8%	1.2%	1.5%	0.6%	1.5%	1.8%	0.9%	0.9%	0.9%	0.6%
More than 6 per day	17.4%	20.7%	6.6%	9.6%	5.1%	5.1%	3.9%	4.5%	5.4%	3.0%	2.1%	2.4%	3.3%

Carrot, Spinach, Broccoli, Cabbage, Peas, beans, onion, garlic, tomatoes, Avocado Tofu, soya meat, and TVP Vege-burger are the least consumed by the participants, never or less than once/per month. Peas, beans, Avocado Tofu, soya meat, and TVP Vege-burger none of the participants consumed more than six times per day (Table 5).

**Table 5** Vegetables consumption per month

	Carrot	Spinach	Broccoli	Cabbage	Peas	Beans	Onion	Garlic	Tomatoes	Avocado	Tofu, soya meat, TVP, Vege-burger
Never or less than once/month	35.4%	48.6%	47.4%	23.1%	43.5%	42.3%	23.4%	25.5%	21.0%	59.5%	71.8%
1-3 per month	22.5%	22.5%	22.8%	18.9%	19.25%	22.8%	21.9%	24.9%	17.7%	14.7%	16.8%
2-4 per week	25.8%	17.1%	16.5%	28.2%	22.5%	21.3%	25.8%	24.0%	31.5%	13.2%	5.4%
5-6 per week	12.6%	9.0%	11.1%	20.7%	12.3%	11.4%	19.2%	17.7%	17.4%	9.9%	3.9%
once per day	3.0%	1.8%	1.5%	7.5%	2.1%	1.8%	5.4%	3.9%	7.8%	2.4%	1.8%
2-3 per day	0.0%	0.3%	0.0%	0.0%	0.0%	0.0%	1.5%	1.5%	2.4%	0.3%	0.3%
4-5 per day	0.3%	0.3%	0.3%	0.6%	0.3%	0.3%	0.9%	0.3%	0.6%	0.0%	0.0%
More than 6 per day	0.3%	0.3%	0.3%	0.9%	0.0%	0.0%	1.8%	2.1%	1.5%	0.0%	0.0%

## 4. DISCUSSION

The medical issue with an inadvertent onset is hypertension, which may cause blood vessels to explode and cause bleeding in organs like the brain or kidneys with fragile capillary beds. Hypertension exceeds obesity, diabetes mellitus, and smoking as the primary risk factors for cardiovascular disease-associated morbidity and mortality (Ahmed et al., 2017). Hypertension and obesity are significant indicators of cardiovascular disease-associated impairment and premature death; hypertension has vast effects on human capital (Chobanian et al., 2003). The prevalence of hypertension and obesity increased with rapid expansion, economic development, acceleration of population aging, and modifications in lifestyle and traditional dietary habits (Geaney et al., 2015). Because it contributes to vascular problems, coronary heart disease (CHD), and stroke, high blood pressure is a condition that should be taken seriously (El-Bcheraoui et al., 2014).

Epidemiological change, an aging population, urbanization, and an increase in the age-specific rates of several chronic disorders are all factors that may be contributing to the total burden of diseases linked with hypertension (Al-Ghamdi et al., 2018). Our study results highlight risk factors for cardiovascular illnesses in the Arar market population are the same as studies done earlier (James et al., 2014). The overall estimated prevalence of hypertension and obesity in our study is comparable to another research (Lu et al., 2022). For female gender, the overweight problem appears to be substantially more prevalent than in earlier research (Mills et al., 2016; Steptoe et al., 2004; Mohammed, 2019; Staessen et al., 2017). Comparatively, the majority of studies Omidi et al., (2021), Adeloye et al., (2014), Alsheikh-Ali et al., (2014) found that men were more likely to have hypertension than women, whereas just a small number of studies Al-Mohaissen et al., (2017) found that women were more likely to have it.

The results of our study provide strong evidence that obesity, overweight problems, and hypertension increase with aging, similar to those found in earlier studies (Mohamed, 2019; Staessen et al., 2017; Omidi et al., 2021). Central obesity is well known to be significantly associated with hypertension (Omidi et al., 2021; Adeloye et al., 2014; Alsheikh-Ali et al., 2014; Al-Mohaissen et al., 2017; Reddy et al., 2004). Many studies accept that central adiposity and hypertension are highly correlated (Reddy et al., 2004).

Those who reported being single in our study had higher blood pressure values than those who had been married. Being single was substantially related to having higher blood pressure levels (Lipowicz et al., 2005; Jeppesen et al., 2000; Khan et al., 2013). Sedentary lifestyle, hypertension, and obesity were the main risk factors for CVD in the market population in our study.

Hypertension and obesity were found to be more common, especially among men in the market population in Arar. In Saudi Arabia, men (18.7%) were more likely than women (14.0%) to have hypertension (Di-Chiara et al., 2017; Whelton et al., 2018). The Eastern province of Saudi Arabia has the most significant prevalence of cardiovascular-associated diseases (Whelton et al., 2018). According to some research, men are more likely than women to have cardiovascular-associated diseases, independent of age (Ferguson et al., 2008; Tsai et al., 2005). There are a few limitations to this search. It is not to extrapolate the findings to the entire market population of Saudi Arabia due to the cross-sectional approach.

Our study was limited to Arar; therefore, it may be difficult to generalize the findings to the entire country. This study is significant and sheds light on the prevalence of risk factors for cardiovascular diseases in Arar, which is significantly high. In our study, the market population with poor diet knowledge was associated with a higher risk factor for CVD. Diet modification can decrease the prevalence of hypertension (Kawamoto et al., 2008; Yu et al., 2008; Darnton et al., 2004). Normotensive and hypertensive adults have shown a marked drop in blood pressure following a healthy diet. A healthy diet encourages consuming more fruit and vegetables while consuming less fat, sodium, and processed foods (Fung et al., 2008).

## 5. CONCLUSION

Overall, the results of the current study reported risk factors for CVD, especially hypertension and obesity, to be prevalent, particularly among adult male sedentary shopkeepers. Awareness campaigns directed towards increasing people knowledge about risk factors for obesity and hypertension are warranted. Awareness programs are needed to educate people about the association of dietary patterns (effects) and physical activity levels with the risk of obesity and hypertension. Our study results highlight that individuals with excellent nutrition and diet quality knowledge are more likely to consume healthier diets, which may significantly influence dietary behaviors among the market population, helping to reduce the incidence of cardiovascular diseases like hypertension, heart attack, and stroke.

### Acknowledgements

The authors sincerely thank the research committee for granting us approval for data collection.

### Author's Contributions

The authors confirm their contribution to the paper as follows: Study conception and design: 1. Anshoo, Khalid; data collection: 1. Abdulaziz, 2. Abdulaziz; analysis and interpretation of results: 1. Amin, 2. Rahma; draft manuscript preparation: 1. Eslam, 2. Anshoo. All authors reviewed the results and approved the final version of the manuscript.

### Ethical Approval

The study received ethical approval from the Northern Border University-Local Bioethics Committee with decision number (12/44/H).

### Informed Consent

Not applicable.

### Funding

This study has not received any external funding.

### Conflict of interest

The authors declare that there is no conflict of interests.

### Data and materials availability

All data sets collected during this study are available upon reasonable request from the corresponding author.

## REFERENCES AND NOTES

- Adeloye D, Basquill C. Estimating the prevalence and awareness rates of hypertension in Africa: a systematic analysis. *PLoS One* 2014; 9(8):e104300. doi: 10.1371/journal.pone.0104300
- Ahmed AM, Hersi A, Mashhoud W, Arafah MR, Abreu PC, Al-Rowaily MA, Al-Mallah MH. Cardiovascular risk factors burden in Saudi Arabia: The Africa Middle East Cardiovascular Epidemiological (ACE) study. *J Saudi Heart Assoc* 2017; 29(4):235-243. doi: 10.1016/j.jsha.2017.03.004
- Al-Ghamdi S, Shubair MM, Aldiab A, Al-Zahrani JM, Aldossari KK, Househ M, Nooruddin S, Razzak HA, El-Metwally A. Prevalence of overweight and obesity based on the body mass index; a cross-sectional study in Alkharj, Saudi Arabia. *Lipids Health Dis* 2018; 17(1):134. doi: 10.1186/s12944-018-0778-5
- Al-Hamdan N, Saeed A, Kutbi A, Choudhry AJ, Nooh R. Characteristics, risk factors, and treatment practices of known adult hypertensive patients in Saudi Arabia. *Int J Hypertens* 2011; 2010:168739. doi: 10.4061/2010/168739
- Al-Mohaissen MA, Al-Obaid QY, AlGhamdi WA, Al-Alyani HS, Dahman SM, Al-Wahhabi NA, Al-Awaji NM, Lee T. Prevalence and Determinants of Hypertension and High Normal Blood Pressure Among Young Adult Saudi Females: Insights in to Preventive Strategies. *Fam Community Health* 2017; 40(3):221-230. doi: 10.1097/fch.000000000000154
- Al-Nozha MM, Ali MS, Osman AK. Arterial hypertension in Saudi Arabia. *Ann Saudi Med* 1997; 17(2):170-4. doi: 10.514/4/0256-4947.1997.170
- Alsheikh-Ali AA, Omar MI, Raal FJ, Rashed W, Hamoui O, Kane A, Alami M, Abreu P, Mashhoud WM. Cardiovascular risk factor burden in Africa and the Middle East: the Africa Middle East Cardiovascular Epidemiological (ACE) study. *PLoS One* 2014; 9(8):e102830. doi: 10.1371/journal.pone.0102830
- Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Edward J. Seventh report of the Joint National Committee on Prevention, detection, evaluation, and treatment of high blood pressure. *Hypertension* 2003; 42(6):1206-1252. doi: 10.1161/01.hyp.0000107251.49515.c2
- Darnton-Hill I, Nishida C, James W. A life course approach to diet, nutrition and the Prevention of Chronic Diseases. *Public Health Nutr* 2004; 7(1A):101-21. doi: 10.1079/phn2003584
- Di-Chiara T, Scaglione A, Corrao S, Argano C, Pinto A, Scaglione R. Education and hypertension: Impact on global cardiovascular risk. *Acta Cardiol* 2017; 72(5):507-513. doi: 10.1080/00015385.2017.1297626
- El-Bcheraoui C, Memish ZA, Tuffaha M, Daoud F, Robinson M, Jaber S, Mikhitarian S, Al-Saeedi M, AlMazroa MA, Mokdad AH, Al-Rabeeh AA. Hypertension and its associated risk factors in the Kingdom of Saudi Arabia, 2013: A national survey. *Int J Hypertens* 2014; 2014:564679. doi: 10.1155/2014/564679
- Ferguson TS, Younger NO, Tulloch-Reid MK, Wright MB, Ward EM, Ashley DE, Wilks RJ. Prevalence of prehypertension and its relationship to risk factors for cardiovascular disease in Jamaica: Analysis from a cross-sectional survey. *BMC Cardiovasc Disord* 2008; 8:20. doi: 10.1186/1471-2261-8-20
- Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L, Alexander L, Estep K, Hassen-Abate K, Akinyemiju TF, Ali R, Alvis-Guzman N, Azzopardi P, Banerjee A, Barnighausen T, Basu A, Bekele T, Bennett DA, Biadgilign S, Catalá-López F, Feigin VL, Fernandes JC, Fischer F, Gebru AA, Gona P, Gupta R, Hankey GJ, Jonas JB, Judd SE, Khang YH, Khosravi A, Kim YJ, Kimokoti RW, Kokubo Y, Kolte D, Lopez A, Lotufo PA, Malekzadeh R, Melaku YA, Mensah GA, Misganaw A, Mokdad AH, Moran AE, Nawaz H, Neal B, Ngalesoni FN, Ohkubo T, Pourmalek F, Rafay A, Rai RK, Rojas-Rueda D, Sampson UK, Santos IS, Sawhney M, Schutte AE, Sepanlou SG, Shifa GT, Shiue I, Tedla BA, Thrift AG, Tonelli M, Truelsen T, Tsilimparis N, Ukwaja KN, Uthman OA, Vasankari T, Venketasubramanian N, Vlassov VV, Vos T, Westerman R, Yan LL, Yano Y, Yonemoto N, Zaki ME, Murray CJ. Global burden of hypertension and systolic blood pressure of at least 110 to 115 mm hg, 1990-2015. *JAMA* 2017; 317(2):165-182. doi: 10.1001/jama.2016.19043
- Fung TT, Chiuve SE, McCullough ML, Rexrode KM, Logroscino G, Hu FB. Adherence to a dash-style diet and risk of coronary heart disease and stroke in women. *Arch Intern Med* 2008; 168(7):713-20. doi: 10.1001/archinte.168.7.713
- Geaney F, Fitzgerald S, Harrington JM, Kelly C, Greiner BA, Perry IJ. Nutrition knowledge, diet quality and hypertension in a working population. *Prev Med Rep* 2015; 2:105-13. doi: 10.1016/j.pmedr.2014.11.008
- James PA, Oparil S, Carter BL, Cushman WC, Dennison-Himmelfarb C, Handler J, Lackland DT, LeFevre ML, MacKenzie TD, Ogedegbe O, Smith SC Jr, Svetkey LP, Taler SJ, Townsend RR, Wright JT Jr, Narva AS, Ortiz E. 2014 evidence-based guideline for the management of high blood pressure in adults: report from the panel members appointed to the Eighth Joint National Committee (JNC 8). *JAMA* 2014; 311(5):507-20. doi: 10.1001/jama.2013.284427

17. Jeppesen J, Hein HO, Suadicani P, Gyntelberg F. High triglycerides and low HDL cholesterol and blood pressure and risk of ischemic heart disease. *Hypertension* 2000; 36(2): 226–32. doi: 10.1161/01.hyp.36.2.226
18. Kawamoto R, Kohara K, Tabara Y, Miki T. High prevalence of prehypertension is associated with the increased body mass index in community-dwelling Japanese. *Tohoku J Exp Med* 2008; 216(4):353–61. doi: 10.1620/tjem.216.353
19. Khan RJ, Stewart CP, Christian P, Schulze KJ, Wu L, LeClerq SC, Khatry SK, West KP Jr. A cross-sectional study of the prevalence and risk factors for hypertension in rural Nepali women. *BMC Public Health* 2013; 13:55. doi: 10.1186/1471-2458-13-55
20. Lipowicz A, Lopuszanska M. Marital differences in blood pressure and the risk of hypertension among Polish men. *Eur J Epidemiol* 2005; 20(5):421–7. doi: 10.1007/s10654-005-1752-x
21. Lu Y, Lan T. Global, regional, and national burden of hypertensive heart disease during 1990–2019: An analysis of the global burden of disease study 2019. *BMC Public Health* 2022; 22(1):841. doi: 10.1186/s12889-022-13271-0
22. Memish ZA, Jaber S, Mokdad AH, AlMazroa MA, Murray CJL, Al-Rabeeah AA; Saudi Burden of Disease Collaborators. Burden of disease, injuries, and risk factors in the Kingdom of Saudi Arabia, 1990–2010. *Prev Chronic Dis* 2014; 11:E169. doi: 10.5888/pcd11.140176
23. Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, Chen J, He J. Global Disparities of Hypertension Prevalence and Control: A Systematic Analysis of Population-Based Studies From 90 Countries. *Circulation* 2016; 134(6):441–50. doi: 10.1161/circulationaha.115.018912
24. Mohamed NA. Prevalence of risk factors for diabetes mellitus and hypertension among adult in Tabuk - Kingdom of Saudi Arabia. *Open Access Maced J Med Sci* 2019; 7(5):831–837. doi: 10.3889/oamjms.2019.046
25. Ng FL, Warren HR, Caulfield MJ. Hypertension genomics and cardiovascular prevention. *Ann Transl Med* 2018; 6(15): 291. doi: 10.21037/atm.2018.06.34
26. Omid N, Arabloo J, Rezapour A, Alaeddini F, Bragazzi NL, Pourasghari H, Behzadifar M, Salarifar M, Khorgami M, Ghorashi SM, Azari S. Burden of hypertensive heart disease in Iran during 1990–2017: Findings from the global burden of disease study 2017. *PLoS One* 2021; 16(9):e0257617. doi: 10.1371/journal.pone.0257617
27. Padmanabhan S, Aman A, Dominiczak AF. Recent Findings in the Genetics of Blood Pressure: How to Apply in Practice or Is a Moonshot Required? *Curr Hypertens Rep* 2018; 20(6):54. doi: 10.1007/s11906-018-0863-1
28. Reddy KS, Katan MB. Diet, nutrition and the prevention of hypertension and cardiovascular diseases. *Public Health Nutr* 2004; 7(1A):167–86. doi: 10.1079/phn2003587
29. Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Makuc DM, Marcus GM, Marelli A, Matchar DB, Moy CS, Mozaffarian D, Mussolino ME, Nichol G, Paynter NP, Soliman EZ, Sorlie PD, Sotoodehnia N, Turan TN, Virani SS, Wong ND, Woo D, Turner MB; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Executive summary: Heart disease and stroke statistics—2012 update: a report from the American Heart Association. *Circulation* 2012; 125(1):188–97. doi: 10.1161/cir.0b013e3182456d46
30. Roy A, Praveen PA, Amarchand R, Ramakrishnan L, Gupta R, Kondal D, Singh K, Sharma M, Shukla DK, Tandon N, Reddy KS, Krishnan A, Prabhakaran D. Changes in hypertension prevalence, awareness, treatment and control rates over 20 years in National Capital Region of India: Results from a repeat cross-sectional study. *BMJ Open* 2017; 7(7):e015639. doi: 10.1136/bmjopen-2016-015639
31. Saeed AA, Al-Hamdan NA, Bahnassy AA, Abdalla AM, Abbas MA, Abuzaid LZ. Prevalence, Awareness, Treatment, and Control of Hypertension among Saudi Adult Population: A National Survey. *Int J Hypertens* 2011; 2011:174135. doi: 10.4061/2011/174135
32. Staessen JA, Li Y, Hara A, Asayama K, Dolan E, O'Brien E. Blood Pressure Measurement Anno 2016. *Am J Hypertens* 2017; 30(5):453–463. doi: 10.1093/ajh/hpw148
33. Steptoe A, Willemsen G. The influence of low job control on ambulatory blood pressure and perceived stress over the working day in men and women from the Whitehall II cohort. *J Hypertens* 2004; 22(5):915–20. doi: 10.1097/00004872-200405000-00012
34. Tsai PS, Ke TL, Huang CJ, Tsai JC, Chen PL, Wang SY, Shyu YK. Prevalence and determinants of prehypertension status in the Taiwanese general population. *J Hypertens* 2005; 23(7):1355–60. doi: 10.1097/01.hjh.0000173517.68234.c3
35. Whelton PK, Carey RM, Aronow WS, Casey DE Jr, Collins KJ, Dennison Himmelfarb C, DePalma SM, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ, Muntner P, Ovbigele B, Smith SC Jr, Spencer CC, Stafford RS, Taler SJ, Thomas RJ, Williams KA Sr, Williamson JD, Wright JT Jr. 2017ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task

- Force on Clinical Practice Guidelines. *Hypertension* 2018; 71 (6):1269-1324. doi: 10.1161/hyp.0000000000000066
36. Yu D, Huang J, Hu D, Chen J, Cao J, Li J, Gu D. Prevalence and risk factors of prehypertension among Chinese adults. *J Cardiovasc Pharmacol* 2008; 52(4):363–8. doi: 10.1097/fjc.0b013e31818953ac